

ASHEESH GUPTA, MD, MPH

www.asheeshguptamd.com

www.novaorthospine.com

Arthroscopic Hip Surgery Physical Therapy Protocol

The intent of this protocol is to provide guidelines for your patient's therapy progression. It is not intended to serve as a recipe for treatment. We request that the PT/PTA/ATC should use appropriate clinical decision making skills when progressing a patient forward.

Please contact Dr. Gupta's office to obtain the operative report from our office prior to the first post-op visit. Also please contact Dr. Gupta if there are any questions about the protocol or your patient's condition.

Please keep in mind common problems that may arise following hip arthroscopy: hip flexor tendonitis, adductor tendonitis, sciatica/piriformis syndrome, ilialupslips and rotations, LB pain from QL hypertonicity and segmental vertebral rotational lesions. If you encounter any of these problems please evaluate, assess, and treat as you feel appropriate, maintaining Dr. Gupta's precautions and guidelines at all times. Gradual progression is essential to avoid flare-ups. If a flare-up occurs, back off with therapeutic exercises until it subsides.

Please reference the exercise progression sheet for timelines and use the following precautions during your treatments. Thank you for progressing all patients appropriately and please send all progress notes to Dr. Gupta; office or hand deliver with the patient themselves. **Successful treatment requires a team approach, and the PT/PTA/ATC is a critical part of the team!** Please contact Dr. Gupta at any time with your input on how to improve the therapy protocol.

Please Use Appropriate Clinical Judgment During All Treatment Progressions

Initial Pre-op Assessment

Assess bilateral hips

ROM – flexion, extension, IR, ER, abd, add

Gait – look for Trendelenburg gait

Impingement test – flexion/adduction/IR often reproduces pain

Ober test

Strength – abduction, flexion, extension

****PLEASE SEE LAST PAGE FOR MODIFICATIONS - PATIENT SPECIFIC PROCEDURES****

Begin therapy POD #1 (unless otherwise instructed)

NoVa Orthopedics and Spine Care
14605 Potomac Branch Drive, Suite 300
Woodbridge, VA 22191
P: (703) 490-1112 F: (703) 8788735

ASHEESH GUPTA, MD, MPH

www.asheeshguptamd.com

www.novaorthospine.com

Phase 1- Immediate Rehabilitation (1-2 weeks):

Goals:

Protection of the repaired tissue
Prevent muscular inhibition and gait abnormalities
Diminish pain and inflammation

Precautions:

20 lb. flat-foot weight-bearing post-op, duration per Dr. Gupta's orders depending on procedure
Do not push through pain or pinching, gentle stretching will gain more ROM

Gentle PROM only, **no passive stretching**

Avoid Capsular Mobilizations

Avoid any isolated contraction of iliopsoas

Initial Exercises

AAROM: within range limitations, pain free.

ROM Guidelines (pain free)

Flexion: 90° x 3 weeks

Ext: 0° x 3 weeks

Abd: 25-30° x 3 weeks

IR: 90 deg. hip flexion: 0deg x 3 weeks; neutral (prone): within comfort zone

ER: 90 deg. hip flexion: 30 deg x 3 weeks; neutral (prone): 20 deg x 3 weeks

After 3 weeks, gradually progress ROM as tolerated, **within pain-free zone*

- STM (scar, anterior, lateral, medial and posterior aspects of hip, lumbar paraspinals, quad/hamstring)

-Stationary bike with no resistance

-Isometric (quad setting, gluteal setting, TA isometrics with diaphragmatic breathing)

-Prone lying (modify if having low back pain) – AVOID in instability patients

Phase 2 – Intermediate Rehabilitation (3-8 weeks)

Criteria for progression to Phase 2:

Full Weight Bearing Must Be Achieved Prior To Progressing To Phase 2

Non weight bearing exercise progression may be allowed if patient is not progressed by MD to full weight bearing (Please see last page for microfracture modifications)

Goals:

Protection of the repaired tissue
Restore Full Hip ROM – (ROM must come before strengthening)
Restore Normal Gait Pattern
Progressive Strengthening of Hip, Pelvis, and LE's
Emphasize gluteus medius strengthening (non-weight bearing)

Precautions:

No forced (aggressive) stretching of any muscles
No joint/capsular mobilizations – to avoid stress on repaired tissue
Avoid inflammation of hip flexor, adductor, abductor, or piriformis

Intermediate Exercises

Gentle strengthening; ROM must come before strengthening

-Stationary bike no resistance, add resistance at 5-6 weeks

NoVa Orthopedics and Spine Care
14605 Potomac Branch Drive, Suite 300
Woodbridge, VA 22191
P: (703) 490-1112 F: (703) 8788735

ASHEESH GUPTA, MD, MPH

www.asheeshguptamd.com

www.novaorthospine.com

- Hooklying progression: pelvic clock, TA w bent knee small range ER, marching, add isometric w kegel ball, isometric abduction with ring.
- Prone progression: IR/ER AROM, prone on elbows with glut setting-press ups, hip extension, alternating arm/leg raise.
- Sidelying progression: clams 30 deg hip flexion to 60 deg hip flexion, hip abduction straight leg raise, side plank on elbow.
- 1/2 kneel: gentle pelvic tilt for gentle stretch of iliopsoas
- Bridge progression
- Balance progression: double leg to single leg balance
- Pelvic floor strengthening
- Elliptical / stair stepper: 6-8 weeks
- Step and squat progression
- Slide board: hip abduction / adduction, extension, IR/ER. No forced abduction. Stop short of any painful barriers.
- Continue to avoid any isolated contraction of iliopsoas

Phase 3 – Advanced Rehabilitation (9-12 weeks)

Criteria for progression to Phase 3:

- Full ROM
- Pain free Normal gait pattern
- Hip flexor strength of 4/5
- Hip abd, add, ext, and IR/ER strength of 4+/5

Goals:

- Full Restoration of muscular strength and endurance
- Full Restoration of patient's cardiovascular endurance
- Emphasize gluteus medius strengthening in weight bearing**

Precautions:

- No contact activities
- No forced (aggressive) stretching
- No joint mobilizations – to avoid stress on repaired tissue

Exercises:

- No treadmill walking until 12 weeks
- 4-pt lumbar / core stabilization progression
- Anterior / side plank progression
- Crab / monster walk
- Lunges all directions
- Single leg squat
- Continue progressions of exercises in phase II.

Phase 4 – Sport Specific Training >12 weeks

Criteria for progression to Sport Specific Training:

- Hip flexor strength 4+/5
- Hip add, abd, ext, IR/ER 5-/5
- Cardiovascular endurance equal to pre-injury level
- Demonstrates proper squat form and pelvic stability with initial agility drills, stable single-leg squat.
- Return to sport activities as tolerated without pain, consistent with MD orders.

NoVa Orthopedics and Spine Care
14605 Potomac Branch Drive, Suite 300
Woodbridge, VA 22191
P: (703) 490-1112 F: (703) 8788735

ASHEESH GUPTA, MD, MPH

www.asheeshguptamd.com

www.novaorthospine.com

Exercises:

- Customize strengthening and flexibility program based on patient's sport and/or work activities
- Z cuts, W cuts, Cariocas
- Agility drills
- Jogging
- Gradual return to sport

NoVa Orthopedics and Spine Care
14605 Potomac Branch Drive, Suite 300
Woodbridge, VA 22191
P: (703) 490-1112 F: (703) 8788735

ASHEESH GUPTA, MD, MPH

www.asheeshguptamd.com

www.novaorthospine.com

MODIFICATIONS for SPECIFIC Procedures

See operative report for specifics and consider the following therapeutic techniques.

Iliopsoas Release:

Begin gentle stretch beginning with prone lying (Phase 1)

Gentle active release of iliopsoas (Phase 2)

Microfracture:

20 lbs FFWB with crutches x 8 weeks

Can progress from Phase 1 to non-weight bearing strengthening portions of Phase 2

Begin full weight bearing at 8 weeks

Capsular Plication for Hip Laxity:

Avoid combined Extension and External Rotation for 6 weeks

Focus rehab on gradual strength progression

No joint mobilizations or over stretching ROM

No Prone ROM for 6 weeks

Gradually progress AAROM under patient's control within comfort

Piriformis Release:

POD #1 begin stretch piriformis (flexion, adduction, ER) without causing anterior hip pain and sciatic nerve flossing (Phase 1)

Gentle active release of piriformis (Phase 2)

Recommended sitting position when having to sit for longer durations.

(Right leg in picture is surgical leg)



NoVa Orthopedics and Spine Care
14605 Potomac Branch Drive, Suite 300
Woodbridge, VA 22191
P: (703) 490-1112 F: (703) 8788735

ASHEESH GUPTA, MD, MPH

www.asheeshguptamd.com

www.novaorthospine.com

Examples of Strengthening



Heel Squeeze

Sitting in a chair with good posture and feet well supported, turn your heels in towards one another so that they are touching (toes/knees apart) and gently squeeze together. Hold 10 seconds. Come back to the center so that toes are pointing straight ahead and relax 10 seconds. Add theraband to progress.

Toe Squeeze

Sitting in a chair with good posture and feet well supported, turn your toes and knees in towards one another so that they are touching (heels apart) and gently squeeze toes together. Hold 10 seconds. Come back to the center (so that toes are pointing straight ahead) and relax 10 seconds. Progress using ball between knees.



-Bridge. Gently squeeze your buttocks as you raise your hips off the ground, knees bent and feet shoulder width apart.

-Bridge (knees together and feet apart)

-Bridge (knees apart and feet together)

NoVa Orthopedics and Spine Care
14605 Potomac Branch Drive, Suite 300
Woodbridge, VA 22191
P: (703) 490-1112 F: (703) 8788735

ASHEESH GUPTA, MD, MPH

www.asheeshguptamd.com

www.novaorthospine.com



Slideboard Exercises (Small range
avoiding hip discomfort)
-Abduction / Adduction
-Extension
-Internal/External Rotation
-Twist

NoVa Orthopedics and Spine Care
14605 Potomac Branch Drive, Suite 300
Woodbridge, VA 22191
P: (703) 490-1112 F: (703) 8788735