ASHEESH GUPTA, MD, MPH www.asheeshguptamd.com www.novaorthospine.com

Dr. Gupta's Hip Medication Instructions

1.	Pain Medication:	
		Norco 10/325mg- take every 6 hours as needed for pain This is a narcotic pain medication. Do NOT take with alcohol or while operating a motor vehicle. Do NOT combine with Tylenol or anything containing acetaminophen
		Other:
2.	Blood Thinners- Anticoagulation:	
		ASA 325mg- take 1 tablet twice a day for 30 days. Finish this medication This is NOT to be taken in patients under 18 years of age
		Other:
3.	Anti-inflammatory- Heterotopic ossification prevention:	
		EC-Naprosyn 500mg- take 1 tablet twice a day for 30 days. Finish this medication Do NOT combine with other anti-inflammatory medications
		Other:
4.	Digestive Medications:	
		Protonix 40mg- take 1 tablet once a day as needed for acid reflux
		Colace 100mg- take 1 tablet up to three times per day as needed for post-operative digestion and constipation
		Zofran 4mg- take as needed for nausea and vomiting every 4-6 hours
		Other:
5.	Antibiotic (only needed if outpatient total hip replacement):	
		Keflex 500mg- take 1 tablet every 6 hours. Start day after surgery. Finish this medication
	П	Other: